An Essay

Compression of the Mrains; submitted to the Commination of the Revent I Andrews: D. D. Mount;

The Thusteen

And Medical Faculty of the University of Neurosylvania,

x Hypoc whitly it as se multin

mus (fo

Eing me

toy it,

the order

The intricacy of this disorder, has engaged the attention of many writers from the day, of Hypocrates sown to the present time: and although some valuable improvements have been made, by modern Surgeons, in the treat = ment of it; yet it appeares still to be im -: perfectly understood. However, I have selected it as the subject of this egray, not with a view (for I have not the presumption to suppose myself capable) of advancing any thing new, or elucidatory of what has been so ably investigated before; but my object is, by connecting the various authors who have treated of it, companing bacir different opinions &c. 5 to become an well arguainted as I can with a Ineadful aggection to which man is so

paetures Ru con to lost ato ever best of aternal Estinguis liable, and with which every practitioner must expect frequently to meet.

Compression of the brain may be produced by various courses; such as, Fractures attended with a depregned portion of the cramium, Extravasortion, and the forma = - tion of matter. It may also be coursed by Hydrocepaulus internus, or by a trickning of bone from Diseaso, as in the lues veneres. The two last mentioned course will not be taken into comberation; but I shall endeavour to treat of here theree first, as occurring from external violence, by bescribing their symptoms, Entinguishing them from other affections apportantly similar, and indicating the means most commonly employed in the cure.

3th of compression of the brain from Depression Fractures.

which o ale the In the Eigenent Past it very begin but that thy wer

The Symptom are stupor, trouving, lop of speed and voluntary motion, humorthage from the ears and non, sicking at the Stomach succeeded by vomiting. There symptom, or a part of been at least, when produced by the above mentioned come, occur immediately after the accident; which circumstances distinguishes this from ale the other course of compregion. In the Heatment of Fractures, authors are of Liggerent opinion, and non more exceptionable Then the justey celebrated Both. He afrests that it is not only indispensably necessary that every depression of the cramium should be elevated but that we should operate in Seven cases out of ten in simple undepressed fractures: but I believe there are very few of the present

fay who would not object to this doctrine.

meani eght to to com y speries a a pres melief & hufeon ly Mu

I should suppose that it would be necessary to in no case, whatever, in consequence of as fracture whether depression existed or not; meggin there were obvious symptoms of a compregned brain, or unless it was evident from the gratured points irretating the Dura Mater inglamation might superview. It is certained right to raise every degreefeed piece of boyo, if it can be some without perforating Instru ments or much violence. But Nov. Noth is of opinion that a perforation should be made as a preventativo of ofuturo mischief: What unselvet can it prevent? We Inglamation.

If Inflamation occur, it is the effect of viocclines or force down to the head; and certain by the application of Instruments connot diminuch this cause but must rather tend por to me F, in to Ter, by ut a se in ou e K Hous & mony by the as bone in with Lynefrion gi

tens to moreouse it. It bylamotion exist, butter ments counst remove it. voly may me not attempt to prevent it, or if it occurs to resour it, in this part of the System on in any other, by copious Brood letting, Junging, Hentering and a strict antephlogistid regimen? It is two it is on a vitale organ, and for that reason we revoued be continue of precipitately operating. I have seen several instances, and read of many others, where a portion of the cran-: im has been depressed by external violence as by the kick of a hono and the patient has some well without its being clevated, and even without the out of a Surgeon. But why a degrapion of the cramin should not at all g times give exemptions of compression, since its eavity in health is completely fileed, I am not

- have win con nat in mupnegn :ranuum " incli when to account, unless on Doctor Huyrick has rug : poset it probable, that the prefoure is made on the longitudinal sinus. We are told that somtimes only a portion of ble internal teches of the Skues is depresent. It would be well to have this fact in view in performing an operation. But I presume no becisivo progmore could be formed of it, untice the first take were perforated. His, however, could not be mate viate in betermining on an operation; for I am! persenaled that when the symptoms of a compressed brain one Distinctly marked, whether produced by bone or any other course, we show never heritate with regard to perforating the cramium: The mode of doing which, by the "repline will hereafter be described. Though I am inclined to think that the Sam invented by In they will be found preferable in most cases

the egg brain. avey the metures, at grown The or ng on ra housed · broin miny from synefred fraction, especially when a would be necessary to apply two mepline

more them oneo.

1 way of compression of the brain grow Estrawasa tion. This Estrawasation of blood or rerum may be the effect of oughtures befores, either between the Cramium and Sura Mater, Na Sura & pria maters, the pia mater & the brain, or in the ventricles of the brain. The Symptoms of Extravaration one warry the same on those occurring from depressed Tractures, with which it is often accompanies, and from which it may be destinguished by not occurring immediately I suddenly, but gradually with our intervalt, generally, between the time of the accident and the time of compressions long or short in proportions to the siges of the Superior refrels. Estravarations is also often accompanied with concussion or commetions of the brown. When the consugation is stight or of

whility. In stee 1 multer at state mengrove confesse part suration, sa aportient generally recover his somes, and after a while, according to been degree of effusions, again relayers into a state of insen sibility. But when the consulpions is considerate. the brain much disordered, I believe their is no greater desideratum in Surgery blan some do invo mark whereby to sertinguish it from compressions. We part of see professions appears better culculated to embarage & perpes vero judgement of the merperienced Surgeon them this In the various authors that I have auxiously committed on this point, I am sorry to say I have found nothing positive or betterments; they sile with one voice asknowledge it dark and obstrue. Mr. I Bell, by rediculing were term concussion & reasoning analogically, has endeav ared to court some light on it, but it must be confessed his vous are founts & glimmering.

2 syrugi courties winn, very; leg * ensing lat ben 7 aut coes, it serve Junes ca "he sies" to says, There is every reason to believe consugation to be a sufferior into the substances of the brain escenbling the ecclymosis of a bruse limb, attended with symptoms resembling apoplesy & terminating, mally in a graduer obserption and rear recove = my; soutines in light inflamatory action & subten belevium, in Exploramies: While compression, proeleding from Estravoration of blood, is plainly maley; begining in Strepor, without insensibility not ending in humplegia & convulsion! He also of terves in another place, when speaking of consugation in that New Fortient, When lifted from the ground, is found cold, pale, motivules, without pulse, without sense and is as in a fainting git; if he continues cold & pages his unine & feces involuntaine ly, he dies"; out goes on to say if he does male his state will be succeeded by high arterial actions recompanies with belevium, which last are good

w often weion . 4 estrem wheneo ! pregin

signs. One when speaking of compression from quesion of blood, sorys, "The "partient is never insenwible unless he is about to expire and namely blesion; his pulse never vises but is slow, heavy spouring at every fifth stroke; his pupie is ometimes belated, one rise usually paralised, the , ther open shakes with a sort of tremulous con welsion, which return at requear intervals; & attended with a quivering irrequeer pulse, set extremities & insensiber evacuations when in is shout to expire"! I common say from aperience whether the Distinction from by Iw. Belo ever good in practice, I fear they to not; if they so very certainly revould have not influence in directing sen more of prace ties. Some cuttion regret bleat we have at "an infallible whereby to destinguish what a profing ferrid is! This in a practical point

1 hours 5 ascerta my to me to Estrai a extern nt. Her mation Mebolomy on the mir. N gent, it the dees o to one

y view can be of no importance. When the symptoms of compressed brain one truly marked, " I have before soid, the only indication then, to assertain the seat of preguire, and endea vous to remove it. The deat of prefrue we are ournary led to by our injured sealy or grature but Extravorsation sometimes takes places without my externor mark of violence to insiente its: teat. Here again we are in the bask. In this. extraction Mr. Both, this' a bold operator, has untry confesses, "the only chance of releif is from no valebotomy and on open belley! There together with jurgeet quietuses & low diet rhould be carried as for on the state of the system will possibly whenite. Mr. Cooper observes of the symptoms and ingent, it certainly might be proper to perforate in cramium in the course of the spinous artery ; for sura mater. If no blood should be found wher one parietal bows, the operation might be

. He wer the Der funct last t nably & ster her i are de bu theres the gene 1. Home to tuere

me on the other! This Mr. Aberrethey's observations so to coroberate; he is of opinion that serious surequences will relative result "rules one of the large exteries of the Dura Matter be wown po" He also mentions of them be so much blood in the dura Mater on materially to desaurge to functions of the brain," he believes "the four so situated will not be found to blees. This last circumstances, presume, would measurably depend on the length of time the Duss mater has been reperated from the hour. we are directed by Mr. Belo, in difficult cares. where there is no external most to direct, to wate general prepure over the heart, and by This, possibly, some tensor spot may be disesser it. However if we can, by any means betermin on the situation of the prefuse, we should ben summediately operate, in blu mouner here after described: our after a sufficiency of bone is

He ops 1 sugar not be within at beris which

surver, if the fluir be found on the Dura mater it may generally be removed, here symptoms releive , & the patient may so well. But if the ferris be between the sura & pria mater the can is extrervely hazarrous. This may always be suspectively tet, when the sura mater is found to protouted at the perforation, giving a sense of tension to ver gingen, and her symptom not mitegated. W. Foll here recommends that the Sena Mater should be punctured; out also the prie mater if meeliary, that is, if the course of prefuse be situa ter within it. But boctor "Mayrick is of opinion that this more of practice is lightly improper: in every case where he has seen The burn mater punctured, Inflamation & beath have been the uniform consequences; and therefore he prefers the meral remisies, particularly copious Bloodletting by which treatment he has known reverae to be

menfrior Wi right all the line heremands that the bone heater H, our laces he purationed, and also her pro male I would go source up the course of most mitently buiso, i it to pro Sympton in indie myle in - the he

I now come to speak of the terrs and last proposed come of Compression of the brain, viz. For Forma tion of Matter within the Cramium. This is generally the consequence of Inflormation; rais Inflormation may be produced by continion. concussion on Fracture. In order to understand bein sightly Mr. Note tells it is necessary to know that the vegues of the perieranium, thou of the diploo, of the dura mater within to all constantly & freely communicates with each other, and upon this greedom of communication depends the health of the parts. Therefore whatever tends to bruiso, jinetato, on sisonder blu circulation, must tend to produce impermotion, unless prevented by the means, commonly employed for this purpose. The Symptoms, I presume, are generally remitar to thon indicating Inflormation of any ether parts, though in a more viveent begree; ruch as hair in the head on if bound by a cost, restlegues.

merion Fem. Fe ag from - accider mation un soon who, and kinas lost; but want of seep, hard & frequent pulse, hot & dry skin, fewhet face, infermed eyes, nausies, vomiting rigors (sign of suppuration) and toward the end convuesion & belesium. This cause of com - prefriow may be distinguished from all other courses of compression, from consugaion or com= motion by see long interval between see time of the accident of the occurrence of the begung = town. The Sepurptoms of a compressed brain, aris ing from other causes, or those of consuguion appear instanteneously or in a short time after the accident; whereas those arising from the Formation of Matter from Inflamation selfour occur sooner Man several Days, sometimes several weeks, and in a few cases in several months. It was the general opinion of former surgeous, . that the matter or his found on the brain or its membranes, mas originally effected or extravorat blood; but this is obviously incomet, as has been

light of some there & traduct from the A look Leffeet. 400 eq with all may believe with he and respon writego tile The w, bee " heart roumens a strict would ! that D. " opera stinfactority proved by Mr. Nott and others. Enfection of the membranes or brain is often an effect of contunion. The first symptoms and, pain in the part which received the blow, which is soon extended over the head, attended with lan your or a dejection of sperits, which is soon for lowed by a mausie and on melination to vomit, vertigo or gibbings, a quick that pulso, and an incapacity of sleeping at least quietty. while the Matient is in this conditions, if the man titioner were to confine him to a dark & quiet toom, been copiously (five or six times a day if the state of his system would admit shows the heart and Breister its over, publish last is stronger recommended by Boets Hayrick) and confino him to a strict antepheogistic regimen; I am confident In would selsow find limself compelled to resort to short disagreeable, Sungerous & less alternative the operation. But if this pean he now pursued

w & rig 4 Taker

the Symptom must go on to mercaro; the part wjured wice become turniques, sope, tenser &pappy; if the Perieranium be examined, to will be found seperated from see cranium and see home of a whitish evenir, here wound discharging, in stead of pur, beloody serum or a bark coloured ichor. In teris situation Mr. Foto would say apply the Frequences without hesitation, remove a portion of how, and you will find the sura mater detachet from the skull and a quantity of matter (the come of compression) situated either upon or within it. This admonition would be put & right; for no soult see process of suppure has taken peace, matter has formed and is forming and unless vent is made for its discharge, the Symptom, to use the words of the learned outloor every hour after this period are examperated, and advance with harty strides; the headach & thirst become more intense, the though becreases, the

a many the state of the state o The year good a live servered from her commen I what is notice eating too mount buckinging in given out of few bloody server or a last convers um ye clare be their tetrestion are not now warden naterio we its womo & & birch Vierouri m die ca year to Murrape ya ma

region are more frequent and at last convulsive motions, attended in some with belevium, in other with paralysis or comators stupicity, finish van tragely. In the above I have spoken eleigtey of short Infermation & suppuration occurring from continion; but from whatever course they may proceed the gebrilo symptom we not materially defferent. If the scale be wounded or cut, though appeared to be doing were will have its aspect considerably changed upon the recurrence of fever. The edges of the some will become heard, painful, pale, loon its granulating, and discharge only a tern discolvered somies; the resievanium will be found, also, seperated & the home discoloused. There two last mentioned symptoms uppear to have determined Mr. Note, not only withrespect to bee propriety of operating, but also withrespect to the detached space of the hura water, I portion of bone to be removed.

1 conec and I be made were at me and plagant and stably worton there there of date by samistion 8 200 was from the work out of the contract of the c witte: m. 141 is of ferre. The odge of the prope muisi ace is & I mi in treator & Karledon on Lane. Ho he w

If concerption be the cause of Informaction, the first Engagement generally broughear, and the Patient appears to enjoy health; but in a gen day, or probably weeks, according to circumstances. The Enjoyetom, of Information Succeed, which grow this came about all others through the resolved if population; for here, generally speaking, it is not ong more extensive, but we have no quido whitever to conduct in to the resit of matter when formed. Information is not unfrequently produced by a combination of course, but her her her here

Information is not emprogrently produced by a combination of cause; but let it be produced by a transition are the same by what is to record it if populate; But if in spite y remibies suppuration currents and the compreparation is evident; as next numb be immediately made, if there he a spot to marrout the appeirant of the most of soing this I mile now endeavour to describe.

July to 4 pregue John News regury. at This lu mue

Trephining. This should rank among spira : tion of the first importance in Surgery; through safe and simple in the lands of a skieful and operienced operator, yet it is one on which the yound & inespesienced connot too much referet. to is one that requires judgement & skill, not only to know when it should be performed, which is of the first moment, but also how to perform it. and it is one "Fact the removal of prefeure off see brain, which prefeure must also actually occasion dangerous symptoms; com form see only true & vindicable reason for per -forming it. However, when from the various symptoms given before, it is judger indespensally meegrang, it should not be delayed. and the fine things to be provided are the Sustruments. Here much might be said withrespect to the surprisonatings & exceety of those used by the autients;

withere sale pro He in . hetter ; Hunge De Bone thouses Meon Myerry

and withrespect to the propriety or impropriety of wany described by blu moderns. But as "Practition in of see present day appear, pritty generally, to withingurs to those that should be used, and as a knowledge of their form, sign, figure tel. may be best obtained by sever semonotration, I hade proceed to describe blu Operation on tought in this university. The furtraments to be potter in reading are 1st a Scalpel provided with a steel edge set the end of New hunder; 2nd The improved Fregerino or two of the same wife would be better; 300 cm Elevator; 4th a Toothpick; 5th a small breest; and 6th Tenaculum, needles, ligatures & springs. The Batients head tip it has not been done) should now be showed, or at least so unch of the hair should be removed, as meespary. be oracies then be laid on a Table with his head on pileon securet & supported by africtants. The seat & injury should now again be examined. and then

nu courte mest en ion the as as their form try desire to i may mgle 1 w ven movet. he m whitely lit dire etent. hitinguis tenly as

no

" one contion of high importance to the mape vienced surgeons, that is in examining a tumor be is very liable to be decrived by a semation of fracture when in fact there is now. upon the nature of the can depend the next step; this may be easily ascertained, if the duteguments happen to be lacerated & see cramium exposed; ix not, an Incipion should be made with the Scalpe through the contusion or Tumor down to the bono; quarting at the same time, against a grow tuno, least be knife might slip thro' and in jum the brain. No part of the scalp should be removet, a simple or crucial insignow is all that can be meegrany. Only hemorrhage must be immediately stoped by prefune or ligature. If a Frantise be found, the breignow should be continued in it direction if it is judged necessary to discover its extent. It has been mentioned by some as defficient is distinguish a Fracture from a duturo; but this may be easily assertained, first by an areatomical knowledge

here blee whe & amino and see " the species desired to company to the many by the we part to in tu Min. put sece mailton bether bear materially to form in bristance the ment your steam

0.

Surverage of the parts; recorder, by a Fraction being ways rough, & thirdey by the Beneramin wohering, imply to a Setture, but not to a Frenchese. When the Fracture is accompanies with depression it should be raised by the Elevator; if this can not be some, the judgement of the operator must setermino whether the saw be preferable to the Superino; in most of term cares I think it will a found preferable, especially where it is judged weegsary to remove a long, narrow or large portion y bone. If the preference be given to the Trepline a perieranium services be seraped off with the hour tu of New Scalpel, and New center pin fixed on a bound part of bone, but so as for the circle of the four to include a portion of vere Fracture. The witer him, being movable, at first should protouso beyond the teeth of the saw, I be naised gradually " the "perforation is going on, untile the Trepline become steady, then it should be finally removed. the grove should often be examined with the hothpick,

tonly to which se In he bu get, wit was not you to travely don't get a species and got on the out he us Mai y also & be cute les blue " Heliet o of the Bealgue, and the center his first on a Those ex mation In being 4 to me

0.

it only to know when it was marrythro, but also a which sito to bear the Saw. Buring this time, if were he but one herr, an agrittent should be enrages, with the Brush or Tower, in cleaning its teeth. when the Saw is found to be nearly terrough, it would be better for fear of its teethe injuring the tura mater, to break out the piece of how with & Elevator; the splinters of bow if then be any way selso be removed by the same instruments. ome cutton mention the deplot as a quide for knowing when the Saw is money through, but this should not a relies on; for many parts of the cranium, especies by teron of old persons have no diplow. after the information is made the define for portion of bone may commonly be mained by introducing one end of the Elevator and bearing on the other. If the depression " accompanies with extravaration it will probably he duchanges this ver perforation.

wan then is no practice, and the operation is performed to relieve progress from blood, wromen or matter towards be conducted in the came manners un.

& my a from to know when the was many true, but also a which siso to hear the form Herring this trine is tiere he hut one how you agritant should be en court with the Bruk or Fourt, in cleaning its test of you were her Saw is found to be nearly tenouse it all a vous he better for fear as its teste injuries the is one was moter, to break out the fire of how will . my a leverton the openter of hour of tour he sum and it was also be grounded by his same anythouse of top our suffer mustion bey biples on a quite for the is it is now they form in regard, through, here saw himselve thatt. trade of the property from the legion. Las ano · but. 1 Vor Kus 小石研

alternative

for often the Brephino should be appeied, she circumstorner of the can and the good sense of the Gurgeon can alone betermine; It is certainly right to remove o much how and only so much as will permit the were of prefoure to be discharged. After the removal of bone, if we find the course ap refour cituated upon the Dura Mater, we conclude ans feer it is favourable; it may be discharged aro' the perforation, and the symptoms of a com prefect brain releived. But on the contrary, if the use be situated under on within the Sura mater, we met hat, refeet Aproceed no farther; but nother y what virtue there is in general remidies, net as must copious bloodletting, lasitives, blistens on diet, rest \$0. Nurrenny is, also, recommended by vetor Rush in despieuet cares, the good effects of while he has witnessed in the Benagenania Hospital. at when the symptoms are extremely ungent, the dura later much protruded, its evenur active 40. I am dis med to think, it would be well as the last

Epilitic City otto Report the Suno Mister, I was con mo: Ou the Younge horshag theeker Washier the war agraphy was a way of a recommendation of the same wheter who by Ranger conation, to punction it with the point of a lancet out when reduced to this extremity, the Patient is tray - property; sure me may now with propriety say to course est be his, a quebus non availant aliquis in north deligion.

Much has been said with respect to operating on continuen parts of the cramium; but I believe suns yeour up her present day relson historic to aprevato in any part that may require it. However in quanting on bone of imquear thickness the surgeon avues be particular in frequently examining the moone; and when the some has entered at on part boots Whyrich, in order to defend the Dura matter, has who suces, with advantage, a small spatials. umorrhage from any of the vegices may generally a cheeked by preging with the finger for some time " introducing a small bit of lint, or when the cirting embetet in bone a small cetar plug as receon mended by that's by sick.

with regard to drefing the wound after the

the seal ton Young 21000, 46 to bevo beging the in a he ture " my 9 aut oliv the en

meration most Bractitioner recommend the geogra , the scalp to be laid down, and a little rope lint weres with a simple outment to be applied; but sector Vayrick objects here use of lint for this good wow, that the line generally become enemented itte blood or some femil & is very deficult to more without much irretating ble parts; and weefore he prefers a craft positive which can easily . removed & renewed as accasion may require. a drefrings are best retained by a common night for the wound is bright the Prosetthouser attention should in be turned particularity to great against duplama time end Lymptomatic gever. In order to expect this, his atient should be committed to a back & quiet soons; infinit to a low diet, and his bonces kept regular gentle levitives. But if symptoms of fever, unforunately, repervent, we must immediately gley to sew bureet as our chief oil; in no case probably is hastetting so coming naquired. This together with beiten

1. more alalogue) In Mint the You empt rent! Bu in whise thater and her welf agentic plan must be strilly

The draw I treated the subject; and altho I have spoken of externor violence as the remove course, yet compression, no soulet frequency occurs from within cause; and after opposes in the form of the most formitable discours in the wordsque Entalogued & buscid such is the sympathy existing between the organ (the brain) and the other parts of the byslem Mat in no discour should it escape the attention of the Pronetitioner. Its functions 80: sine, downers, at present so imperfectly known, that it would be present prior in me to say any string faither, I show There for concerns choping, with Boston Cullen, This weeat. But befor I conclude I humbey tender my contract Ranks to the mediene Franciscon, executively & interior heally, for the much information I have obtained from his lectures, and went they may long live to occupy the wais which they now no ably file is the fewent wish of their serverter Gervert.

